arXiv:1412.3310v1 [math.DG] 10 Dec 2014

Legendre surfaces with harmonic mean curvature vector field in the unit 5-sphere *

Toru Sasahara

Abstract

We obtain the explicit representation of Legendre surfaces in the unit 5-sphere with harmonic mean curvature vector field, under the condition that the mean curvature function is constant along a certain special direction.

2000 *Mathematics Subject Classification* Primary 53C42; Secondary 53B25

1 Introduction

It is well-known that an odd-dimensional unit sphere S^{2n+1} is equipped with the standard Sasakian structure (g, ϕ, η, ξ) (see, Section 2). The study of minimal Legendre submanifolds in S^{2n+1} is a very active field and closely related to one of minimal Lagrangian submanifolds in complex projective space.

A natural generalization of a minimal submanifold is a submanifold with parallel mean curvature vector field. However, Legendre submanifolds in S^{2n+1} with parallel mean curvature vector field are in fact minimal (see, [11]). So, as a generalization of minimal Legendre submanifolds, it is natural to consider Legendre submanifolds whose mean curvature vector field H is harmonic with respect to the normal Laplacian Δ^D (see, Section 2), that is,

$$\Delta^D H = 0. \tag{1.1}$$

The purpose of this paper is to study the class of nonminimal Legendre surfaces satisfying (1.1) in the unit 5-sphere. The mean curvature function $||H||$ of such surfaces is not constant (see, Lemma 2 in Section 3). From this, the class seems to be very large. Thus, we need to add some natural condition. On nonminimal Legendre submanifolds, there exists a special vector field: ϕH. Moreover in case the dimension is 2, up to signs, there is a unique unit vector field normal to ϕH. We denote it by $(\phi H)^{\perp}$. In this paper, under the condition that the square of the mean curvature function is constant along one of ϕH and $(\phi H)^{\perp}$, we completely determine nonminimal Legendre surfaces satisfying (1.1) in the unit 5-sphere as follows:

Theorem 1. *Let $f : M^2 \to S^5 \subset \mathbf{C}^3$ be a nonminimal Legendre surface satisfying $\Delta^D H = 0$ in the unit 5-sphere. Then the mean curvature function of M^2 is not constant. If H satisfies $\phi H||H||^2 = 0$, then f is locally given by*

$$f(x,y) = \left(\frac{1}{\sqrt{2}} \exp(\frac{1+\sqrt{5}}{2} iy) \cos x, \frac{1}{\sqrt{2}} \exp(\frac{1-\sqrt{5}}{2} iy) \cos x, \sin x \right). \tag{1.2}$$

*Rocky Mountain Journal of Mathematics **40** (2010), 313-320. An appendix is added.

If H satisfies $(\phi H)^{\perp}||H||^2 = 0$, then f is locally given by

$$f(x,y) = \frac{1}{\sqrt{2}}\left(i + \sin x, (\sec x + \tan x)^i \cos x \cos y, (\sec x + \tan x)^i \cos x \sin y\right). \tag{1.3}$$

2 Legendre submanifolds in the unit sphere

Let $\mathbf{C}^{n+1}$ be the complex Euclidean $(n+1)$-space together with the canonical complex structure J. Denote by S^{2n+1} the unit sphere with the standard induced metric g in $\mathbf{C}^{n+1}$. The position vector field $\mathbf{x}$ of S^{2n+1} is a unit normal vector field of S^{2n+1} in $\mathbf{C}^{n+1}$ and the vector field $\xi := -J\mathbf{x}$ is tangent to S^{2n+1}. Define a 1-form η and an endomorphism field φ on M by the formula:

$$JX = \phi X + \eta(X)\mathbf{x}, X \in TM.$$

It is easy to see that (g, φ, η, ξ) satisfies

$$\varphi^2 = -I + \eta \otimes \xi,\ d\eta(X,Y) = g(X, \varphi Y).$$

Thus (g, φ, η, ξ) is a contact metric structure of S^{2n+1}.

Denote by $\overline{\nabla}$ the Levi-Civita connection of S^{2n+1}. Then

$$\overline{\nabla}_X \xi = -\varphi X,\ (\overline{\nabla}_X \varphi)Y = g(X,Y)\xi - \eta(Y)X.$$

These formulas imply that S^{2n+1} is a Sasakian manifold.

An immersed n-submanifold $x : M^n \to S^{2n+1}$ is said to be a *Legendre submanifold* if $x^*\eta = 0$. The formulas of Gauss and Weingarten of x are given respectively by

$$\begin{aligned} \bar{\nabla}_X Y &= \nabla_X Y + h(X,Y), \\ \bar{\nabla}_X V &= -A_V X + D_X V, \end{aligned} \tag{2.1}$$

where $X, Y \in TM^n$, $V \in T^{\perp}M^n$, ∇, h, A and D are the Levi-Civita connection of M^n, the second fundamental form, the shape operator and the normal connection. The mean curvature vector H is given by $H = \frac{1}{n}\text{trace}h$. Its length $||H||$ is called the *mean curvature function* of M^n. The normal Laplacian is defined by $\Delta^D = -\sum_{i=1}^n (D_{e_i} D_{e_i} - D_{\nabla_{e_i} e_i})$, where $\{e_i\}$ is a local orthonormal frame of M^n.

For Legendre submanifolds we have [1]

$$A_{\phi Y} X = -\phi h(X,Y) = A_{\phi X} Y, \quad A_\xi = 0. \tag{2.2}$$

Moreover a straightforward computation shows that the equations of Gauss, Codazzi, Ricci of Legendre submanifolds in the unit sphere are equivalent to

$$\langle R(X,Y)Z, W\rangle = \langle [A_{\phi Z}, A_{\phi W}]X, Y\rangle + \langle X, W\rangle \langle Y, Z\rangle - \langle X, Z\rangle \langle Y, W\rangle, \tag{2.3}$$

$$(\bar{\nabla}_X h)(Y,Z) = (\bar{\nabla}_Y h)(X,Z), \tag{2.4}$$

where $\bar{\nabla} h$ is defined by $(\bar{\nabla}_X h)(Y,Z) = D_X h(Y,Z) - h(\nabla_X Y, Z) - h(Y, \nabla_X Z)$.

3 The proof of Theorem 1

We assume that the mean curvature function is nowhere zero. Let $\{e_i\}$ $(i = 1, \ldots, 5)$ be an orthonormal frame along M^2 such that e_1, e_2 are tangent to M^2, $\phi e_1 = e_3$, $\phi e_2 = e_4$, $\xi = e_5$ and $H = \frac{\alpha}{2}\phi e_1$, with $\alpha > 0$. Then, it follows from (2.1) and (2.2) that the second fundamental form takes the form:

$$\begin{aligned} h(e_1, e_1) &= (\alpha - c)\phi e_1 + b\phi e_2, \\ h(e_1, e_2) &= b\phi e_1 + c\phi e_2, \\ h(e_2, e_2) &= c\phi e_1 - b\phi e_2, \end{aligned} \tag{3.1}$$

for some functions b, c.

We put $\omega_i^j(e_k) = \langle \nabla_{e_k} e_i, e_j \rangle$. From (2.4) we get

$$e_1 c + 3b\omega_1^2(e_1) = e_2 b + (\alpha - 3c)\omega_1^2(e_2), \tag{3.2}$$

$$-e_1 b + 3c\omega_1^2(e_1) = e_2 c + 3b\omega_1^2(e_2), \tag{3.3}$$

$$e_2(\alpha - c) - 3b\omega_1^2(e_2) = e_1 b + (\alpha - 3c)\omega_1^2(e_1). \tag{3.4}$$

Suppose that M^2 satisfies $\Delta^D H = 0$. Then we have the following three lemmas (see p.290 and 291 in [10]).

Lemma 2.

$$\Delta_M \alpha + \alpha\{1 + (\omega_1^2(e_1))^2 + (\omega_1^2(e_2))^2\} = 0, \tag{3.5}$$

$$2(e_1\alpha)\omega_1^2(e_1) + 2(e_2\alpha)\omega_1^2(e_2) + \alpha\{e_1(\omega_1^2(e_1)) + e_2(\omega_1^2(e_2))\} = 0, \tag{3.6}$$

$$e_1\alpha + \alpha\omega_1^2(e_2) = 0, \tag{3.7}$$

where Δ_M is the Laplace operator acting on $C^\infty(M)$.

Note that α is not constant by (3.5), since α is nowhere zero.

Lemma 3. *There exists local coordinates x, y such that*

$$\begin{aligned} e_1 &= \alpha\partial_x, \\ e_2 &= \alpha\partial_y, \\ \omega_1^2(e_1) &= \alpha_y, \\ \omega_1^2(e_2) &= -\alpha_x. \end{aligned}$$

Lemma 4. *The following relation holds:*

$$b^2 = \frac{\alpha c}{2} - c^2. \tag{3.8}$$

The allied mean curvature vector $a(H)$ is defined by $\sum_{r=4}^{5}(\mathrm{trace}A_H A_{e_r})e_r$ (see, p.197 in [2]). If $a(H)$ vanishes identically on M^2, it is called a *Chen* surface.

Suppose that M^2 is not Chen surface, i.e., $b \neq 0$. Then we may assume that $b > 0$, if necessary by changing the sign of e_2. By differentiating (3.8) we get

$$b_i = \frac{(\alpha - 4c)c_i + \alpha_i c}{4b}, \tag{3.9}$$

where $i = x, y$.

Using (3.9) we replace (3.2) and (3.3) by the derivatives with respect to x and y as follows:

$$\begin{pmatrix} \alpha & -\dfrac{\alpha(\alpha-4c)}{4b} \\ \dfrac{\alpha(\alpha-4c)}{4b} & \alpha \end{pmatrix} \begin{pmatrix} c_x \\ c_y \end{pmatrix} = \begin{pmatrix} -(\alpha-3c)\alpha_x - \dfrac{12b^2-\alpha c}{4b}\alpha_y \\ \dfrac{12b^2-\alpha c}{4b}\alpha_x + 3c\alpha_y \end{pmatrix}. \tag{3.10}$$

First we investigate the case of $\phi H||H|| = 0$, i.e., $\alpha_x = 0$. Then using (3.8) and (3.10) we get

$$c_x = -\frac{8bc}{\alpha^2}\alpha_y, \tag{3.11}$$

$$c_y = \frac{5\alpha c - 8c^2}{\alpha^2}\alpha_y. \tag{3.12}$$

By a long but straightforward computation, we obtain

$$\begin{aligned} bc_{xy} = {} & (-24\alpha_y^2 - 4\alpha\alpha_{yy})\left(\frac{c}{\alpha}\right)^2 + (112\alpha_y^2 + 8\alpha\alpha_{yy})\left(\frac{c}{\alpha}\right)^3 \\ & -128\alpha_y^2\left(\frac{c}{\alpha}\right)^4, \end{aligned} \tag{3.13}$$

$$bc_{yx} = -20\alpha_y^2\left(\frac{c}{\alpha}\right)^2 + 104\alpha_y^2\left(\frac{c}{\alpha}\right)^3 - 128\alpha_y^2\left(\frac{c}{\alpha}\right)^4, \tag{3.14}$$

Since $bc_{xy} = bc_{yx}$, we find that $\alpha_y^2 + \alpha\alpha_{yy} = 0$ or $c = 0$ or $\alpha = 2c$. If $\alpha_y^2 + \alpha\alpha_{yy} = 0$, it contradicts to (3.5). If $c = 0$ or $\alpha = 2c$, it is a contradiction to $b \neq 0$ by (3.8).

Next, we investigate the case of $(\phi H)^\perp||H|| = 0$, i.e., $\alpha_y = 0$. Similarly as in the case of $\alpha_x = 0$, we have

$$c_x = \frac{-3\alpha c + 8c^2}{\alpha^2}\alpha_x, \tag{3.15}$$

$$c_y = \frac{4\alpha b - 8bc}{\alpha^2}\alpha_x, \tag{3.16}$$

$$\begin{aligned} bc_{xy} = {} & -6\alpha_x^2\left(\frac{c}{\alpha}\right) + 56\alpha_x^2\left(\frac{c}{\alpha}\right)^2 \\ & -152\alpha_x^2\left(\frac{c}{\alpha}\right)^3 + 128\alpha_x^2\left(\frac{c}{\alpha}\right)^4, \end{aligned} \tag{3.17}$$

$$\begin{aligned} bc_{yx} = {} & (-4\alpha_x^2 + 2\alpha\alpha_{xx})\left(\frac{c}{\alpha}\right) + (48\alpha_x^2 - 8\alpha\alpha_{xx})\left(\frac{c}{\alpha}\right)^2 \\ & -(144\alpha_x^2 - 8\alpha\alpha_{xx})\left(\frac{c}{\alpha}\right)^3 + 128\alpha_x^2\left(\frac{c}{\alpha}\right)^4, \end{aligned} \tag{3.18}$$

Hence we get $\alpha_x^2 + \alpha\alpha_{xx} = 0$ or $c = 0$ or $\alpha = 2c$, since $bc_{xy} = bc_{yx}$. It is a contradiction as mentioned above.

Therefore we conclude that b must be 0, i.e., M^2 must be a Chen surface if $\phi H||H||^2 = 0$ or $(\phi H)^\perp|H||^2 = 0$. Applying the classification of Legendre Chen surfaces satisfying $\Delta^D H = 0$ (see, Theorem 8 and Corollary 9 in [9]), we can prove the statement.

4 Other examples of Legendre surfaces satisfying $\Delta^D H = 0$

In this section, we show a way to construct Legendre surfaces satisfying $\Delta^D H = 0$ and $(\phi H)^\perp ||H||^2 \neq 0$ and $\phi H ||H||^2 \neq 0$.

One can obtain the following existence and uniqueness theorem by the similar way to those given in [6] and [5] (cf. [7]).

Theorem 5. *Let $(M^n, \langle\cdot,\cdot\rangle)$ be an n-dimensional simply connected Riemannian manifold. Let σ be a symmetric bilinear TM^n-valued form on M^n satisfying*

(1) *$\langle\sigma(X,Y),Z\rangle$ is totally symmetric,*

(2) *$(\nabla\sigma)(X,Y,Z) = \nabla_X\sigma(Y,Z) - \sigma(\nabla_X Y, Z) - \alpha(Y,\nabla_X Z)$ is totally symmetric,*

(3) *$R(X,Y)Z = \langle Y,Z\rangle X - \langle X,Z\rangle Y + \sigma(\sigma(Y,Z),X) - \sigma(\sigma(X,Z),Y)$.*

Then there exists a Legendre isometric immersion $x : (M^n, \langle\cdot,\cdot\rangle) \to S^{2n+1}$ such that the second fundamental form h satisfies $h(X,Y) = \phi\sigma(X,Y)$.

Theorem 6. *Let $x^1, x^2 : M^n \to S^{2n+1}$ be two Legendre isometric immersions of a connected Riemannian n-manifold into the unit sphere S^{2n+1} with second fundamental forms h^1 and h^2. If*

$$\left\langle h^1(X,Y), \phi x^1_* Z\right\rangle = \left\langle h^2(X,Y), \phi x^2_* Z\right\rangle$$

for all vector fields X, Y, Z tangent to M^n, there exists an isometry A of S^{2n+1} such that $x^1 = A\circ x^2$.

Let $f(t)$ be a solution of the following ODE:

$$\frac{d^2 f}{dt^2} = \frac{1}{2}e^{-2f}. \tag{4.1}$$

We put $\alpha(x,y) := e^{f(x-y)}$. Let $(M^2, g = \frac{1}{\alpha^2}(dx^2+dy^2))$ be a Riemannian 2-manifold. We define a symmetric bilinear form σ on M^2 by

$$\begin{aligned}
\sigma(e_1,e_1) &= \frac{3}{4}\alpha e_1 + \frac{\alpha}{4}e_2,\\
\sigma(e_1,e_2) &= \frac{\alpha}{4}e_1 + \frac{\alpha}{4}e_2,\\
\sigma(e_2,e_2) &= \frac{\alpha}{4}e_1 - \frac{\alpha}{4}e_2,
\end{aligned} \tag{4.2}$$

where $e_1 = \alpha\partial_x$ and $e_2 = \alpha\partial_y$. By a straight-forward computation, we find that $((M^2,g),\sigma)$ satisfies (1), (2) and (3) of Theorem 5. Therefore there exists a unique Legendre surfaces in S^5 whose second fundamental form h is given by $h = \phi\sigma$. Moreover such a surface satisfies $\Delta^D H = 0$ and $(\phi H)^\perp ||H||^2 \neq 0$ and $\phi H||H||^2 \neq 0$.

Appendix: Correction to Theorem 1 (Added on December 10, 2014)

As stated previously, Theorem 1 can be proved by applying Theorem 8 and Corollary 9 in [9]. However, Corollary 9 in [9] is incorrect. The immersion (44) in [9] should be replaced by the immersion (4.9) in [4]. Also, the immersion (45) in [9] should be replaced by the immersion (4.6) in [3] such that λ is constant.

Accordingly, (1.2) of this paper should be corrected as follows:

$$f(x,y) = (z(y)\cos x, \sin x)\,, \tag{a.1}$$

where $z(y)$ is a unit speed Legendre curve with nonzero constant curvature in $S^3(1)$.

Also, (1.3) should be corrected as follows:

$$f(x,y) = (z_1(x), z_2(x)\cos y, z_2(x)\sin y), \tag{a.2}$$

where (z_1, z_2) is a unit speed Legendre curve in $S^3(1)$ given by

$$\begin{aligned} z_1 &= (ib + \sin x)/\sqrt{1+b^2}, \\ z_2 &= \exp(ib\ln(\sec x + \tan x))\cos x/\sqrt{1+b^2} \end{aligned} \tag{a.3}$$

for some nonzero real number b.

Remark A.1 (i) A unit speed Legendre curve $z(y)$ of constant curvature λ in $S^3(1)$ is congruent to

$$z(y) = (1/\sqrt{1+b^2})(e^{iby}, be^{-iy/b})$$

with $b = (\lambda + \sqrt{\lambda^2+4})/2$.
(ii) The curvature of Legendre curve given by (a.3) in $S^3(1)$ is $b\sec x$.
(iii) Both surfaces given by (a.1) and (a.2) have constant curvature one.

References

[1] D. E. Blair, *Riemannian geometry of contact and symplectic manifolds,* Progress in Mathematics, Vol 203, Birkhauser Boston, Inc., Boston, MA, 2002.

[2] B. Y. Chen, *A report on submanifolds of finite type,* Soochow J. Math. **22** (1996), 117-337.

[3] B. Y. Chen, *Interaction of Legendre curves and Lagrangian submanifolds*, Israel J. Math. **99** (1997), 69-108.

[4] B. Y. Chen, *Classification of slumbilical submanifolds in complex space forms*, Osaka J. Math. **39** (2002), 23-47.

[5] B. Y. Chen and L. Vrancken, *Existence and uniqueness theorem for slant immersions and its applications,* Result. Math. **31** (1997), 28-39.

[6] B. Y. Chen and L. Vrancken, *Addendum to: Existence and uniqueness theorem for slant immersions and its applications,* Result. Math. **39** (2001), 18-22.

[7] B. Y. Chen, F. Dillen, L. Verstraelen and L. Vrancken, *An exotic totally real minimal immersion of S^3 in CP^3 and its characterization,* Proc. Royal Soc. Edinburgh Ser. A. **126** (1995), 153-165.

[8] J. Inoguchi, *Submanifolds with harmonic mean curvature vector field in contact 3-manifolds,* Colloq. Math. **100** (2004), 163-179.

[9] T. Sasahara, *Legendre surfaces whose mean curvature vectors are eigenvectors of the Laplace operator,* Note. Mat. **22**. (2003), 49-58.

[10] T. Sasahara, *Legendre surfaces in Sasakian space forms whose mean curvature vectors are eigenvectors,* Publ. Math. Debrecen. **67** (2005), 285-303.

[11] K.Yano and M. Kon, *Structures on manifolds*, Series in Pure Mathematics, Vol 3, World Scientific, Singapore, 1984.

Department of System and Information Engineering
Hachinohe Institute of Technology
Hachinohe 031-8501, JAPAN
E-mail address: sasahara@hi-tech.ac.jp